

Cardiac drugs

Beta Blockers

Start with 50% lower dose in patients with eGFR < 30 mL/min/1.73m²

Titrate to effect (monitoring blood pressure and heart rate)

Digoxin

Reduce dose based on plasma concentrations

	eGFR (mL/min/1.73m ²)	
Load	< 15	if loading dose necessary reduce dose by 50%
Maintenance	10–50	administer 25–75% of normal daily dose, or administer Q36H
	< 10	administer 10–25% of the normal daily dose or administer Q48H
Heart Failure <i>intended to target a digoxin concentration of 0.7 ng/mL</i>	30–80	0.125 mg PO daily
	< 30	0.125 mg PO Q48H

Spirolactone

	eGFR (mL/min/1.73m ²)	
Initial dose	30–50	12.5 mg PO daily or every other day
Maintenance	30–50	12.5–25 mg PO daily
	< 30	not recommended

Cardiac drugs

Fenofibrate				
	Normal dose range	eGFR (mL/min/1.73m ²)		
		> 50	20-50	< 20
Fenofibrate micronized	200 mg PO daily; max 200 mg PO daily	100%	Avoid, use alternative formulation	Contraindicated
Lipidil Supra	160 mg PO daily; max 200 mg PO daily	100%	Initiate at 100 mg PO daily	Contraindicated
Lipidil EZ	145 mg PO daily; max 145 mg PO daily	100%	Initiate at 48 mg PO daily	Contraindicated